



Recommended Schedule: [Recommended Immunizations for Children \(Birth through 6 years\) Preteens and Teens](https://www.cdc.gov/vaccines/hcp/vis/index.html)

<https://www.cdc.gov/vaccines/hcp/vis/index.html>

The vaccination recommendations are based on current information and recommendations from the Advisory Committee on Immunization Practice (ACIP) and the American Academy of Pediatrics.

Approximate Age: Vaccine (Combination vaccines will be give when appropriate)

Birth:					
1 month:	Exam Only				
2 month:		*Vaxelis	Prevnar	Rotateq	
4 month:		*Vaxelis	Prevnar	Rotateq	
6 month:		*Vaxelis	Prevnar	Rotateq	
9 month:	Exam Only (if up-to-date on immunizations)				
12 month:	Hepatitis-A	Proquad *	Prevnar		Lead /Hemaglobin
15 month:		Pentacel *			
18 month:	Hepatitis-A			M-Chat	
2 years:	Exam Only (if up-to-date on immunizations)			M-Chat	Lead /Hemaglobin
3 years:	Exam Only (if up-to-date on immunizations)				
4-5 years:		Proquad *	Quadracel *	Vision and Hearing testing	
11 years and up: 2 doses	MenQuadfi	Gardasil	Tdap	Depression Screening 12yr & UP	
		Gardasil (2nd dose/6months after first dose)			
15 years and up will need 3 doses	Gardasil #1	Gardasil #2 1-2 mo from #1		Gardasil #3 6mo from #1 and 3mo from #2	
16 years and up:	MenQuadfi		Trumemba(2nd dose to be given 6mo after the 1st)		

Vaxelis- Hep B; DtaP (Diphtheria, Tetanus, Acellular Pertussis) ; HIB (Haemophilus Influenza Group B) ; IPV (Inactivated Polio)

Prevnar 20 - Pneumococcal Conjugate Valent 20

Rotateq - Rotavirus Vaccine, Live, Oral Pentavalent (Can Not be given after 8 months of age, not even one day older)

Proquad - MMR (Measels, Mumps, Rubella) ; Varicella (Chickenpx Vaccine) (must be 1 year of age)

Hep-A - Hepatitis A (second dose to be given 6 months after the first, no sooner) Recommended

Pentacel - DtaP (Diphtheria, Tetanus, Acellular Pertussis) ; HIB (Haemophilus Influenza Group B) ; IPV (Inactivated Polio)

Quadracel - DtaP (Diphtheria, Tetanus, Acellular Pertussis); Polio (Inactivated Polio)

MenQuadfi - Meningococcal (Groups A, C, Y and W-135 Polysaccharide) 2years and older

Gardasil - Quadrivalent Human Papillomavirus Types 6,11,16,18 Recombinant Recommended

Trumemba - Meningococcal (Serogroup B) 2 doses 6moths apart (recommended for college)

Tdap - Tetanus Toxoids, Diphtheria and Acellular Pertussis Absorbed

Bayfortus (RSV) *Neonates and Infants born during or entering their first RSV season
*Children up to 24 months who remain vulnerable to severe RSV through their 2nd season

COVID (pfizer) 6mo-4y 3 dose series 5yr-11yr 1 dose 12yr-older 1 dose

Influenza - 6 months to 8 years of age should receive 2 doses of vaccine the first year they are vaccinated. 4wk apart

Spot Vision- offer at 6 mo well, if pass repeat at 24 mo then yearly. If the fail the 6 month repeat at 9 month, if the fail repeat at 12 month if they fail send to ophthalmology. Use Snellen chart starting at age 4 years old!

Depression Screening 12 years of age then every year after

Girls 13 years U/A if has menses pregnancy test. 15yr U/A; Preg/GC&C / Boys if sexually active GC&C or if sx

12 years Drug testing for patients on controlled substances